ed ac at 10 moleculor translate a 21 to 24. Each

to Pinet the unit tempent sector, and normal vector and the bisonnal vector at the Unit the Discount Vector at the Unit Pinet Vector and the Unit Pi

podanut edi lo zeulav muminim briti muminim valuez di the function

(7, 0) pure (0, 0)

consulvative or not. If comportative, find its scalar potential function.

is Verific States with later for the verific field  $F(x,y,z) = 2zi + 3zi + 6yi + 6yiz taking the constant of the particle of the <math>x^2 - x^2 + z^2 + z = 0$ , with opward

personance and the universe to be the posterely original circle of intersection of the continued of all of the property of the continued of the property of th



K20U 1535

Reg. No.: .....

Name : .....

V Semester B.Sc. Degree (CBCSS – Reg./Sup./Imp.)
Examination, November 2020
(2014 Admn. Onwards)
CORE COURSE IN MATHEMATICS

Time: 3 Hours Max. Marks: 48

5B08 MAT - Vector Calculus

## SU - SECTION - A LO HOLLO CHIEFTON E DE L'ANDIE CHIEFTON E DE L'AN

All the four questions are compulsory. Each question carries 1 mark :

- 1. Find the gradient of the function  $f(x, y) = tan^{-1} \left(\frac{y}{x}\right)$  at (1, 1).
- 2. If  $w = \sin(x + 2z)$  and  $x^3 + z^3 = 3$ , find  $\frac{dw}{dx}$  using chain rule.
- 3. Evaluate  $\int\limits_{C}9x^{2}y\ dx$  where C is given by  $x=t^{2},\ y=t^{3},\ 0\leq t\leq 2.$
- 4. Write the formula for finding the surface area of a surface S given by F(x, y, z) = C, defined over the planar region R. (4×1=4)

## SECTION - B

Answer any 8 questions from questions 5 to 14. Each question carries 2 marks :

- 5. Write the vector equation of a line in a plane passing through (1, 2) and making an angle  $\frac{\pi}{3}$  with the positive X-axis.
- 6. Find the length of the curve  $r(t) = 3\cos t i 3\sin t j 4t k$ ,  $1 \le t \le 3$ .
- 7. Find the directional derivative of  $f(x, y) = e^{2xy}$  at (-2, 0) in the direction of i + j.

K20U 1535

-2-



- 8. If  $w = x^2 + y^2 + z^2$  and  $z^3 + xyz + yz^2 = 0$ , find  $\frac{\partial w}{\partial x}$  at (x, y, z) = (1, 1, 0) treating x and y as independent variables.
- 9. Find the linear approximation L(x, y) of the function  $f(x, y) = e^{2y+x}$  at (2, 3).
- 10. If  $\phi(x,y) = 3\sqrt{x^2 + y^2}$ , find div(grad( $\phi$ )).
- 11. Find the flux of F = (x y)i + xj across the circle  $x^2 + y^2 = 1$  in the XY-plane.
- 12. If the force F = 4xi + 4yj is acting on a particle moving it along the curve r(t) = ti + (1 + 2t)j from (1, 3) to (3, 7), find the work done by the force.
- 13. Find a parametrization of the surface of the paraboloid  $z = 16 x^2 y^2$ ,  $z \ge 0$ .
- 14. State the Gauss divergence theorem.

(8×2=16)

## SECTION - C

Answer any 4 questions from questions 15 to 20. Each question carries 4 marks :

- Find the equation of the plane through the points A(1, 0, 2), B(1, 1, 1), C(1, 2, 3).
- 16. Show that  $\frac{d}{dt}(U.V) = \frac{d}{dt}(U).V + U.\frac{d}{dt}(V)$ , where U, V are functions of t into  $\mathbb{R}^2$ .
- 17. Use the Taylor's formula for  $f(x, y) = e^x \cos y$  at the origin to find the quadratic approximation of f. Hence find approximate value of f(0.1, 0.2).
- 18. Find Curl (F × G) at (1, 2, 0) where F(x, y, z) =  $3x^2i + 2xyj + 2yzk$  and G(x, y, z) =  $4yzi + y^2j + xyzk$ .
- 19. Using Green's theorem, find the area enclosed by the circle  $x^2 + y^2 = 4$ .
- 20. Evaluate the surface integral  $\iint_{\sigma} y^2 z^2 dS$ , where  $\sigma$  is the part of the cone  $z = \sqrt{x^2 + y^2}$  that lies between the planes z = 1 and z = 2. (4×4=16)



-3-

K20U 1535

## SECTION - D

Answer any 2 questions from questions 21 to 24. Each question carries 6 marks :

- 21. Find the unit tangent vector, unit normal vector and the binormal vector at t = 0 for the curve r(t) = 2cost i + 2sint j + 4tk.
- 22. Find the absolute maximum and minimum values of the function f(x, y) = 3xy 6x 3y + 7 on the closed triangular region R with vertices (0, 0), (3, 0) and (0, 5).
- 23. Check whether the vector field  $F = (6xy + z^3)i + (3x^2 z)j + (3xz^2 y)k$  is conservative or not. If conservative, find its scalar potential function.
- 24. Verify Stoke's theorem for the vector field F(x, y, z) = 2zi + 3xj + 5yk, taking the surface to be the portion of the paraboloid z = 4 x² y² for z ≥ 0, with upward orientation, and the curve to be the positively oriented circle of intersection of the paraboloid with the XY-plane. (2x6=12)