



Reg. No. : .....

Name : .....

**Third Semester M.Sc. Degree (Reg.) Examination, October 2018**  
**MATHEMATICS**  
**(2017 Admn. Onwards)**  
**MAT3C14 : Advanced Real Analysis**

Time : 3 Hours

Max. Marks : 80

**PART - A**

Answer **four** questions from this part. **Each** question carries **4** marks.

1. Give an example of a sequence of functions which converges pointwise but not uniformly.
2. If  $\{f_n\}$  and  $\{g_n\}$  converge uniformly on a set  $E$ , prove that  $\{f_n + g_n\}$  converges uniformly on  $E$ .
3. Consider  $f(x) = \sum_{n=1}^{\infty} \frac{1}{1+n^2x}$ . On what intervals does it fail to converge uniformly?
4. Show that  $e^x$  defined on  $\mathbb{R}^1$  satisfy the relation  $(e^x)' = e^x$ .
5. Define orthogonal system of functions and give an example.

6. Prove that  $\lim_{x \rightarrow 0} \frac{\log(1+x)}{x} = 1$ . (4x4=16)

**PART - B**

Answer **4** questions from this part without omitting **any** Unit. **Each** question carries **16** marks.

**Unit - I**

7. a) If  $\{f_n\}$  is a sequence of continuous function on  $E$ , and if  $f_n \rightarrow f$  uniformly on  $E$ , then show that  $f$  is continuous on  $E$ .
- b) If  $f_n \in \mathcal{R}(\alpha)$  on  $[a, b]$  and if  $f(x) = \sum_{n=1}^{\infty} f_n(x)$  ( $a \leq x \leq b$ ), the series converging uniformly on  $[a, b]$ , then prove that  $\int_a^b f d\alpha = \sum_{n=1}^{\infty} \int_a^b f_n d\alpha$ .

P.T.O.



8. a) Even if  $\{f_n\}$  is a uniformly bounded sequence of continuous functions on a compact set  $E$ , prove that there need not exist a subsequence which converges pointwise on  $E$ .
- b) If  $\{f_n\}$  is a pointwise bounded sequence of complex functions on a countable set  $E$ , then prove that  $\{f_n\}$  has a subsequence  $\{f_{n_k}\}$  such that  $\{f_{n_k}(x)\}$  converges for every  $x \in E$ .
9. State and prove Stone Weierstrass theorem.

### Unit - II

10. a) Suppose  $\sum c_n$  converges. Put  $f(x) = \sum_{n=0}^{\infty} c_n x^n$  ( $-1 < x < 1$ ). Then prove that  $\lim_{x \rightarrow 1} f(x) = \sum_{n=0}^{\infty} c_n$ .
- b) Define analytic functions and give an example.
11. a) Suppose the series  $\sum a_n x^n$  and  $\sum b_n x^n$  converge in the segment  $S = (-R, R)$ . Let  $E$  be the set of all  $x \in S$  at which  $\sum_{n=0}^{\infty} a_n x^n = \sum_{n=0}^{\infty} b_n x^n$ . If  $E$  has a limit point in  $S$ , then prove that  $a_n = b_n$  for  $n = 0, 1, 2, \dots$ . Hence  $\sum_{n=0}^{\infty} a_n x^n = \sum_{n=0}^{\infty} b_n x^n$  for all  $x \in S$ .
- b) Let  $\{\phi_n\}$  be orthonormal on  $[a, b]$ . Let  $s_n(x) = \sum_{m=1}^n c_m \phi_m(x)$  be the  $n^{\text{th}}$  partial sum of the Fourier series of  $f$ , and suppose  $t_n(x) = \sum_{m=1}^n \gamma_m \phi_m(x)$ . Then prove that  $\int_a^b |f - s_n|^2 dx \leq \int_a^b |f - t_n|^2 dx$ , and equality holds if and only if  $\gamma_m = c_m$  ( $m = 1, 2, \dots, n$ ).
12. a) If, for some  $x$ , there are constants  $\delta > 0$  and  $M < \infty$  such that  $|f(x+t) - f(x)| \leq M|t|$  for all  $t \in (-\delta, \delta)$ , then prove that  $\lim_{N \rightarrow \infty} S_N(f; x) = f(x)$ .
- b) If  $f(x) = 0$  for all  $x$  in some segment  $J$ , then prove that  $\lim_{N \rightarrow \infty} S_N(f; x) = 0$  for every  $x \in J$ .
- c) If  $f$  is continuous (with period  $2\pi$ ) and if  $\epsilon > 0$ , then prove there is a trigonometric polynomial  $P$  such that  $|P(x) - f(x)| < \epsilon$  for all real  $x$ .



### Unit - III

13. a) Define the dimension of a vector space and give an example.
- b) Define basis of a vector space.
- c) Let  $r$  be a positive integer. If a vector space  $X$  is spanned by a set of  $r$  vectors, then prove that  $\dim X \leq r$ .
14. a) Suppose  $E$  is an open set in  $\mathbb{R}^n$ ,  $f$  maps  $E$  into  $\mathbb{R}^m$ ,  $f$  is differentiable at  $x_0 \in E$ ,  $g$  maps an open set containing  $f(E)$  into  $\mathbb{R}^k$ , and  $g$  is differentiable at  $f(x_0)$ . Then prove that the mapping  $F$  of  $E$  into  $\mathbb{R}^k$  defined by  $F(x) = g(f(x))$  is differentiable at  $x_0$  and  $F'(x_0) = g'(f(x_0))f'(x_0)$ .
- b) Suppose  $f$  maps a convex open set  $E \subset \mathbb{R}^n$  into  $\mathbb{R}^m$ ,  $f$  is differentiable in  $E$ , and there is a real number  $M$  such that  $\|f'(x)\| \leq M$  for every  $x \in E$ . Then prove that  $|f(b) - f(a)| \leq M|b - a|$  for all  $a \in E, b \in E$ .
15. State and prove implicit function theorem. (4×16=64)