



Reg. No. :

Name :

I Semester M.Sc. Degree (CBCSS – OBE – Regular)
Examination, October 2023
(2023 Admission)
MATHEMATICS
MSMAT 01C04 : Topology

Time : 3 Hours

Max. Marks : 80

PART – A

Answer **any five** questions. **Each** question carries **4** marks.

1. Give a basis for the standard topology on \mathbb{R}^2 .
2. Let Y be a subspace of X and A be a subset of Y . Show that A is closed in Y if and only if it equals the intersection of a closed set of X with Y .
3. Show that limits of sequences are unique in a Hausdorff space.
4. Prove or disprove : Product topology is finer than the box topology.
5. Define a quotient map and give an example.
6. Is every path connected space connected ? Justify your answer. (5×4=20)

PART – B

Answer **any three** questions. **Each** question carries **7** marks.

7. Let $X = \{1, 2, 3, 4, 5\}$ and $\mathcal{T} = \{\emptyset, X, \{2, 5\}, \{2, 3, 4\}, \{2, 3, 4, 5\}, \{1, 2, 5\}, \{3, 4\}, \{2\}\}$
 - a) List the closed subsets of X .
 - b) Determine the closure of the sets $\{2, 4\}$ and $\{1, 3\}$.
 - c) Check whether the space (X, \mathcal{T}) is T_2 .
 - d) Check whether the space (X, \mathcal{T}) is connected.

P.T.O.

K23P 3114

-2-



8. Let

$d_1((x_1, x_2), (y_1, y_2)) = [(x_1 - y_1)^2 + (x_2 - y_2)^2]^{1/2}$; $d_2((x_1, x_2), (y_1, y_2)) = \max\{|x_1 - y_1|, |x_2 - y_2|\}$ and $d_3((x_1, x_2), (y_1, y_2)) = \min\{1, [(x_1 - y_1)^2 + (x_2 - y_2)^2]^{1/2}\}$ and $\mathcal{T}_1, \mathcal{T}_2$ and \mathcal{T}_3 be the corresponding topologies on \mathbb{R}^2 induced by the metrics d_1, d_2 and d_3 respectively. Which of the following options are correct ? Justify your answer.

a) $\mathcal{T}_1 = \mathcal{T}_2 \neq \mathcal{T}_3$

b) $\mathcal{T}_1 \neq \mathcal{T}_2 = \mathcal{T}_3$

c) $\mathcal{T}_1 \neq \mathcal{T}_2 \neq \mathcal{T}_3$

d) $\mathcal{T}_1 = \mathcal{T}_2 = \mathcal{T}_3$

9. Consider the subset $A = \{(x \times \sin \frac{1}{x}) : 0 < x \leq 1\}$ of the plane \mathbb{R}^2 .

a) Find the closure of A in \mathbb{R}^2 .b) Determine the connectedness and path connectedness of closure of A .

10. Which of the following are topological property ?

a) connectedness

b) boundedness

c) pathconnectedness.

11. Determine whether the following subspaces of \mathbb{R} are homeomorphic.

a) $[0, 1)$ and $(0, 1)$ b) \mathbb{Q} and \mathbb{Z} c) $(0, 1)$ and $S^1 \setminus \{(1, 0)\}$ where $S^1 = \{x \times y : x^2 + y^2 = 1\}$ considered as a subspace of \mathbb{R}^2 . (3×7=21)

-3-

K23P 3114

PART – C

Answer **any three** questions. **Each** question carries **13** marks.

12. Let X and Y be two topological spaces.

a) If A is a subspace of X and B is a subspace of Y , prove that the product topology on $A \times B$ is the same as the topology $A \times B$ inherits as the subspace of $X \times Y$.

b) If Π_1 and Π_2 are projections of $X \times Y$ to X and Y respectively, then prove that the collection $\{\Pi_1^{-1}(U) : U \text{ is open in } X\} \cup \{\Pi_2^{-1}(V) : V \text{ is open in } Y\}$ is a sub-basis for the product topology on $X \times Y$.

13. a) Let X and Y be topological spaces and $f : X \rightarrow Y$. Prove that the following are equivalent.

i) f is continuous.ii) For every subset A of X , $f(\overline{A}) \subset \overline{f(A)}$.iii) For every closed set B of Y , the set $f^{-1}(B)$ is closed in X .

b) Let $X = A \cup B$ where A and B are closed in X and $f : A \rightarrow Y$ and $g : B \rightarrow Y$ be continuous. If $f(x) = g(x)$ for every $x \in A \cap B$, then prove that f and g combine to give a continuous function $h : X \rightarrow Y$ defined by $h(x) = f(x)$ if $x \in A$ and $h(x) = g(x)$ if $x \in B$.

14. a) Define order topology and linear continuum.

b) Give an example of a linear continuum.

c) Show that a linear continuum in the order topology is connected.

15. a) Show that \mathbb{R}^n in product topology is metrizable.

b) Is \mathbb{R}^n connected in the product topology ? Give details to support your assertion.

16. a) Prove that the composition of two quotient maps is a quotient map.

b) Let X and Y be topological spaces and $p : X \rightarrow Y$ be a quotient map. Let Z be a topological space and $g : X \rightarrow Z$ be a map that is constant on each set $p^{-1}(\{y\})$. Show that g induces a map $f : Y \rightarrow Z$ such that $f \circ p = g$ and f is a quotient map if and only if g is a quotient map. (3×13=39)