



K24P 0864

Reg. No. :

Name :

**Second Semester M.Sc. Degree (C.B.S.S. – Supple. (One Time Mercy
Chance)/Imp.) Examination, April 2024
(2017 to 2022 Admissions)
MATHEMATICS
MAT 2C 09 : Foundations of Complex Analysis**

Time : 3 Hours

Max. Marks : 80

PART – A

Attempt **any four** questions from this part. **Each** question carries **4** marks :

- Given that γ and σ are closed rectifiable curves having the same initial points. Prove that $n(\gamma + \sigma, a) = n(\gamma, a) + n(\sigma, a)$ for every $a \in \{\gamma\} \cup \{\sigma\}$.
- Let f be analytic on $B(0, 1)$ and suppose $|f(z)| \leq 1$ for $|z| < 1$. Show that $|f'(0)| \leq 1$.
- Does the function $f(z) = z^2 \sin\left(\frac{1}{z}\right)$ has an essential singularity at $z = 0$? Justify your answer.
- Using residue Theorem, prove that $\int_0^\infty \frac{1}{1+x^2} dx = \frac{\pi}{2}$.
- Define the set $C(G, \Omega)$ and show that it is non-empty.
- State the Weierstrass Factorization theorem.

PART – B

Answer **any four** questions from this part without omitting any Unit. **Each** question carries **16** marks :

Unit – I

- a) Prove the following : If G is simply connected and $f : G \rightarrow \mathbb{C}$ is analytic in G then f has a primitive in G .
- b) State and prove The Open Mapping Theorem.

P.T.O.

K24P 0864

-2-



- State and prove the Third Version of Cauchy's Theorem.
- Prove the following : let G be a connected open set and let $f : G \rightarrow \mathbb{C}$ be an analytic function. Then the following conditions are equivalent.
 - $f \equiv 0$;
 - there is a point a in G such that $f^{(n)}(a) = 0$ for each $n \geq 0$;
 - $\{z \in G : f(z) = 0\}$ has a limit point in G .

Unit – II

- a) Show that for $a > 1$, Show that $\int_0^\pi \frac{d\theta}{a + \cos\theta} = \frac{\pi}{\sqrt{a^2 - 1}}$.
- b) State and prove the Residue theorem.
- State and prove the Laurent Series Development.
- Prove the following :
 - If $|a| < 1$ then $\phi_a(z) = \frac{z-a}{1-\bar{a}z}$ is a one-one map of $D = \{z : |z| < 1\}$ on to itself ; the inverse of ϕ_a is ϕ_{-a} . Furthermore, ϕ_a maps ∂D on to ∂D , $\phi_a'(0) = 1 - |a|^2$ and $\phi_a'(a) = (1 - |a|^2)^{-1}$.
 - Let $f(z) = \frac{1}{z(z-1)(z-2)}$; give the Laurent series of $f(z)$ in each of the following annuli :
 - $\text{ann}(0; 0, 1)$,
 - $\text{ann}(0; 1, 2)$,
 - $\text{ann}(0; 2, \infty)$.

Unit – III

- a) Prove the following : If G is open in \mathbb{C} then there is a sequence $\{K_n\}$ of compact subsets of G such that $G = \cup_{n=1}^\infty K_n$. Moreover the sets K_n can be chosen to satisfy the following conditions :
 - $K_n \subset \text{int } K_{n+1}$.
 - $K \subset G$ and K is compact implies $K \subset K_n$ for some n .
 - Every component of $C_\infty - K_n$ contains a component of $C_\infty - G$.
- b) State and prove Hurwitz's theorem.



-3-

K24P 0864

- a) With the usual notations, prove that $|1 - E_p(z)| \leq |z|^{p+1}$ for $|z| \leq 1$ and $p \geq 0$.
- b) Discuss the convergence of the infinite product $\prod_{n=1}^\infty \frac{1}{1+n^p}$ for $p > 0$.
- a) Show that $\prod(1+z_n)$ converges absolutely iff $\sum|z_n|$ converges.
- b) Prove the following : If $\text{Re } z_n > 0$ then the product $\prod z_n$ converges absolutely iff the series $\sum(z_n - 1)$ converges absolutely.
- c) Prove the following : Let $\text{Re } z_n > 0$ for all $n \geq 1$. Then $\prod_{n=1}^\infty z_n$ converges to a non zero number iff the series $\sum_{n=1}^\infty \log z_n$ converges.

